arXiv:1404.3850v1 [math.FA] 15 Apr 2014

# AN IMBEDDING OF FRACTIONAL ORDER SOBOLEV-GRAND LEBESGUE SPACES, with constant evaluation.

**E.Ostrovsky[a], L.Sirota[b]**

[a] Corresponding Author. Department of Mathematics and computer science, Bar-Ilan University, 84105, Ramat Gan, Israel.
E - mail: eugostrovsky@list.ru

[b] Department of Mathematics and computer science. Bar-Ilan University, 84105, Ramat Gan, Israel.

E - mail: sirota3@bezeqint.net

**Abstract**. We extend in this article the classical imbedding theorems for fractional Lebesgue-Sobolev's spaces into the so-called Grand Lebesgue spaces, with sharp constant evaluation.

**Keywords and phrases:** Fractional Sobolev spaces, Aronszajn, Gagliardo and Slobodetskii norms, fractional Laplacian, Sobolev's imbedding, Riesz potential, Grand and ordinary Lebesgue spaces, Sobolev-Grand Lebesgue spaces, Sobolev Derivative Grand Lebesgue spaces, Fourier transform, constants evaluation, upper and low estimates.

2010 Mathematical Subject Classification: Primary 46E35, Secondary 35S30, 35S05.

# 1 Introduction. Notations. Problem Statement.

For the (measurable) numerical function $f : R^n \to R$ the Fourier transform $F[f](\xi) = \hat{f}(\xi)$ is defined as ordinary:

$$\hat{f}(\xi) = F[f](\xi) := \int_{R^n} e^{-ix\cdot\xi} \, f(\xi) \, dx,$$

Hereafter $x \cdot \xi$ denotes the inner (scalar) product of two vectors $x, \ \xi \in R^d$ and $|x|$ is ordinary Euclidean norm of the vector $x : \ |x| = \sqrt{x \cdot x}$.

The Fourier transform $F[f](\xi)$ is correctly defined, e.g. if $f \in \cup_{p \in [1,2]} L_p(R^n)$. The norm of the function $f$ in the Lebesgue, more exactly, Lebesgue-Riesz space $L_p = L_p(R^n), \ p \ge 1$ will be denoted for simplicity $|f|_p$ :

$$|f|_p := \left[ \int_{R^n} |f(x)|^p \, dx \right]^{1/p}.$$

Let $\Delta$ be the Laplacian. The fractional, in general case, power $\sqrt{-\Delta}^s$ may be defined as a pseudo-differential operator through Fourier transform

$$\sqrt{-\Delta}^{s}[f] := F^{-1}(|\cdot|^{s}\ F[f]).$$

The fractional Sobolev's space $W_p^s = W_p^s(R^n)$ consists by definition on all the functions $f : R^n \to R$ with finite norm (more precisely, semi-norm)

$$||f||W_p^s \stackrel{def}{=} |(-\Delta)^{s/2}[f]|_p,\ p \ge 1, \tag{1.1}$$

the Aronszajn-Gagliardo norm; which is equivalent to the Slobodetskii $||\cdot||S_p^s$ semi-norm:

$$||f||S_p^s = ||f||S_p^s(R^n) \stackrel{def}{=} \left[\int_{R^n}\int_{R^n} \frac{|f(x)-f(y)|^p}{|x-y|^{n+sp}}\ dx\ dy\right]^{1/p}. \tag{1.2}$$

More information about these spaces, in particular on the imbedding theorem see in the works [31], [32], [33], [35], [40], [41], [42], [43], [44], [45], [47], [48], [49], [50] etc.

**Remark 1.1.** In the definition (1.2) instead the whole space $R^n$ may be used arbitrary open set $\Omega \subset R^n$. In detail:

$$||f||S_p^s(\Omega) \stackrel{def}{=} \left[\int_{\Omega}\int_{\Omega} \frac{|f(x)-f(y)|^p}{|x-y|^{n+sp}}\ dx\ dy\right]^{1/p}. \tag{1.2a}$$

**Remark 1.2.** In the case when $s < 0$ the fractional Laplace operator $(-\Delta)^{-|s|/2}$ coincides with Riesz potential

$$(-\Delta)^{-|s|/2}[f](x) = I_{|s|}[f](x) = C_R(n,s)\int_{R^n} \frac{f(y)}{|x-y|^{n-|s|}}\ dy.$$

This case was detail investigated in many works, see e.g.[29], [34], [37], [39], [15], [16]. Therefore, we can assume further $s > 0$.

**Remark 1.3.** The "complete" norm in the fractional Sobolev's space may be introduced as follows.

$$||f||V_p^s \stackrel{def}{=} \left[|f|_p^p + |\ (-\Delta)^{s/2}[f]\ |_p^p\right]^{1/p} =$$

$$\left[\int_{R^n} |f(x)|^p\ dx + (||f||W_p^s)^p\right]^{1/p},\ p \ge 1. \tag{1.3}$$

**Remark 1.4.** The fractional Sobolev's spaces are closely related with Besov spaces, see [1], page 330-341.

We will use further the following *sharp* imbedding theorem, see [37], [38], section 4.3. Define the following "constant"

$$K(n,s) := \pi^{s/2}\frac{\Gamma((n-s)/2)}{\Gamma((n+s)/2)}\left\{\frac{\Gamma(n)}{\Gamma(n/2)}\right\}^{s/n}, \tag{1.4}$$

where $\Gamma(\cdot)$ denotes the usually Gamma-function. If

$$0 < s < n,\ 1 < p < n/s,\ u \in C_0^{\infty}(R^n),\ q = pn/(n - sp), \tag{1.5}$$

and $q = pn/(n - sp)$, then

$$|u|_q \le K(n,s)|\sqrt{-\Delta}^s\ u|_p = K(n,s)||u||W_p^s - \tag{1.6}$$

fractional Lebesgue-Sobolev's imbedding theorem.

Note that the conditions (1.5) are also *necessary* for the inequality of a form (1.6) for some constant $K(n,s)$. This assertion may be proved by means of the well-known scaling method, see e.g. [28], [25].

Evidently, the inequality (1.6) holds true for all functions $u = u(x)$ belonging to the *completion* of the space $C_0^{\infty}$ relative the fractional semi-norm $||\cdot||W_p^s$.

**Our aim in this article is to extrapolate the fractional Lebesgue-Sobolev's imbedding theorem (1.6) into the so-called fractional Grand Lebesgu-Sobolev's imbedding spaces, and as a particular case the - into the so-called Exponential Orlicz Spaces (EOS).**

A particular (but weight) case was considered in the previous article [25].

# 2 Grand Lebesgue Spaces and Sobolev-Grand Lebesgue Spaces.

Now we will describe using Grand Lebesgue Spaces (GLS) and Sobolev's Grand Lebesgue Spaces (SGLS).

### 1. Grand Lebesgue Spaces.

We recall in this section for reader conventions some definitions and facts from the theory of GLS spaces.

Recently, see [2], [3], [4], [5], [6], [7],[8], [10], [11], etc. appears the so-called Grand Lebesgue Spaces $GLS = G(\psi) = G\psi = G(\psi; A, B),\ A, B = \mathrm{const}, A \ge 1, A < B \le \infty$, spaces consisting on all the measurable functions $f : R^n \to R$, (or more generally $f : \Omega \to R$) with finite norms

$$||f||G(\psi) \stackrel{def}{=} \sup_{p \in (A,B)} \left[|f|_p/\psi(p)\right]. \tag{2.1}$$

Here $\psi(\cdot)$ is some continuous positive on the *open* interval $(A, B)$ function such that

$$\inf_{p\in(A,B)} \psi(p) > 0,\ \psi(p) = \infty,\ p \notin (A,B).$$

We will denote

$$\mathrm{supp}(\psi) \stackrel{def}{=} (A,B) = \{p : \psi(p) < \infty,\} \tag{2.2}$$

The set of all $\psi$ functions with support $\mathrm{supp}(\psi) = (A,B)$ will be denoted by $\Psi(A,B)$.

This spaces are rearrangement invariant, see [1], and are used, for example, in the theory of probability [7], [10], [11]; theory of Partial Differential Equations [3], [6]; functional analysis [4], [5], [8], [11], [14]; theory of Fourier series [10], theory of martingales [11], mathematical statistics [12], [13]; theory of approximation [19] etc.

Notice that in the case when $\psi(\cdot) \in \Psi(A,B)$, a function $p \to p \cdot \log \psi(p)$ is convex, and $B = \infty$, then the space $G\psi$ coincides with some *exponential* Orlicz space.

Conversely, if $B < \infty$, then the space $G\psi(A,B)$ does not coincides with the classical rearrangement invariant spaces: Orlicz, Lorentz, Marcinkiewicz etc.

**Remark 2.1.** If we define the *degenerate* $\psi_r(p), r = \mathrm{const} \geq 1$ function as follows:

$$\psi_r(p) = \infty,\ p \neq r; \psi_r(r) = 1$$

and agree $C/\infty = 0, C = \mathrm{const} > 0$, then the $G\psi_r(\cdot)$ space coincides with the classical Lebesgue space $L_r$.

**Remark 2.2.** Let $\xi : \Omega \to R$ be some (measurable) function from the set $L(p_1,p_2),\ 1 \leq p_1 < p_2 \leq \infty$. We can introduce the so-called *natural* choice $\psi_\xi(p)$ as as follows:

$$\psi_\xi(p) \stackrel{def}{=} |\xi|_p;\ p \in (p_1,p_2).$$

**2. Sobolev-Grand Lebesgue (SGL) spaces.**

Let $\psi = \psi(p)$ be the function described above. We will say that the function $f : R^d \to R$ belongs to the Sobolev-Grand Lebesgue space $SGL\psi$, iff the following it semi-norm is finite:

$$||u||SGL\psi_s \stackrel{def}{=} \sup_{p\in\mathrm{supp}(\psi)} \left[ \frac{||u||W_p^s}{\psi(p)} \right]. \tag{2.3}$$

This notion (up to equivalence) for the integer values $s$ appeared at first (presumably) in an article [26] (2010); the near definition see in [9] (2013).

**3. Main result.**

**Theorem 2.1.** *Let* $\psi(\cdot) \in \Psi(1, n/s)$ *where* $0 < s < n,\ 1 < p < n/s$. *Define the function*

$$\nu(q) := \psi\left( \frac{qn}{n+qs} \right), \tag{2.4}$$

*so that*

$$\operatorname{supp} \nu(\cdot) = ((n/(n-s),\ \infty).$$

*Let also*

$$u \in C_0^{\infty}(R^n) \cap SGL\psi_s. \tag{2.4}$$

*Proposition:*

$$||u||G\nu \le K(n,s) \cdot ||u||SGL\psi_s, \tag{2.5}$$

*where the constant $K(n,s)$ is the best possible.*

**Proof.** Let $u \in C_0^{\infty}(R^n) \cap SGL\psi_s$; we can and will suppose without loss of generality $||u||SGL\psi_s = 1$. It follows from direct definition of Sobolev-Grand Lebesgue spaces

$$||u||W_p^s \le \psi(p).$$

It follows from inequality (1.6)

$$|u|_q \le K(n,s)||u||W_p^s \le K(n,s)\psi(p). \tag{2.6}$$

Since $p = qn/(n+qs)$, we deduce from (2.6) for the values $q > n/(n-s)$

$$|u|_q \le K(n,s)\psi(qn/(n+qs)) = K(n,s)\nu(q) = K(n,s)\ \nu(q)\ ||u||SGL\psi_s, \tag{2.7}$$

or equally

$$||u||G\nu \le K(n,s)\ ||u||SGL\psi_s,$$

The *exactness* of the constant $K(n,s)$ in (2.5) follows immediately from the one of main results, namely, theorem 2.1, of an article [27].

Note that in the case when $\psi(p) = \psi_r(p),\ r = \mathrm{const} \ge 1$ we obtain as a particular case the ordinary fractional Sobolev's imbedding theorem.

# 3 Boundedness of fractional Laplacian in DGLS

Let us return to the fractional Sobolev's inequality (1.6):

$$|u|_q \le K(n,s)|\sqrt{-\Delta}^s\ u|_p = K(n,s)||u||W_p^s. \tag{1.6}$$

Assume now that the inequality (1.6) is true for some *interval* of values $s:\ s \in (s_-, s_+),\ 0 < s_- < s_+ < 1$.

More detail, let $Q = Q(s_-, s_+)$ be some (measurable) set in the plane $(p, s),\ p \ge 1,\ s \in (0,1),\ Q = \{(p,s)\}$ such that

$$\forall s \in (s_-, s_+)\ \Rightarrow \exists p \ge 1,\ u \in W_p^s. \tag{3.1}$$

If for some values $(p,s)\ u \notin W_p^s$, we denote formally $||u||W_p^s = \infty$.

Denote

$$Q_s = \{p,\ p \ge 1,\ (p,s) \in Q\};$$

the "section" of the set $Q$ on the $s-$ level. Then $\forall s \in (s_-, s_+)\ \Rightarrow Q_s \ne \emptyset$.

The inequality (1.6) may be rewritten as follows:

$$|u|_q \le K(n,s)\ ||u||W^s_{qn/(n+qs)},\ s \in (s_-, s_+),$$

therefore

$$|u|_q \le \inf_{s \in (s_-, s_+)} \left[ K(n,s)\ ||u||W^s_{qn/(n+qs)} \right]. \tag{3.2}$$

The last inequality be reformulated on the language of Grand Lebesgue spaces as follows. Denote

$$\zeta(q) = \inf_{s \in (s_-, s_+)} \left[ K(n,s)\ ||u||W^s_{qn/(n+qs)} \right],\ q \in (n/(n-s_+), \infty),$$

then $||u||G\zeta \le 1$.

**Definition of Derivative Grand Lebesgue spaces.**

Let $\tau = \tau(p,s),\ p > 1,\ s \in (s_-, s_+)$ be continuous function such that $\inf_{p,s} \tau(p,s) = 1$. By definition, the function $u = u(x),\ x \in R^n$ (or $x \in \Omega$) belongs to the Derivative Grand Lebesgue space $DGL(\tau)$, if it has a finite semi - norm

$$|||u|||DGL\tau := \sup_{p>1} \sup_{s \in (s_-, s_+)} \left[ \frac{||u||W_p^s}{\tau(p,s)} \right]. \tag{3.3}$$

We can now formulate the imbedding theorems in Derivative Grand Lebesgue spaces.

**Theorem 3.1.** *Let $u(\cdot) \in DGL(\tau)$; then*

$$|u|_q \le \inf_{s \in (s_-, s_+)} [\ K(n,s)\ \tau(qn/(n+qs), s)\ ]\ \cdot |||u|||DGL\tau. \tag{3.4}$$

**Proof** is alike one in the theorem 2.1. Indeed, let $|||u|||DGL\tau = 1$; then

$$||u||W_p^s \le \tau(p,s),\ p = qn/(n+qs).$$

It remains to use the inequality (3.2) and take the minimum over $s$.

**Corollary 3.1.** The inequality (3.4) may be reformulated on the language of GL spaces as follows. Denote

$$\lambda(q) = \inf_{s \in (s_-, s_+)} [\ K(n,s)\ \tau(qn/(n+qs), s)\ ]\,;$$

then

$$|u|_q \le \lambda(q) \cdot |||u|||DGL\tau,$$

or equally

$$||u||G\lambda \le |||u|||DGL\tau. \tag{3.5}$$

# 4 Weight generalization

Let $\Omega$ be an open *convex* subset of a whole space $R^n$. Introduce after R. L. Frank and R. Seiringer [33] (case $\Omega = R^n_+$) and M. Loss and C. Sloane [39] (general case) the following functions, measures and operators:

$$d_\alpha(x) := \inf_{y \notin \Omega} |x-y|^\alpha,$$

$$D_{\alpha,n}(p) := 2\pi^{(n-1)/2}\ \frac{\Gamma((1+\alpha)/2)}{\Gamma((n+\alpha)/2)}\ \int_0^1 \frac{\left|1 - r^{(\alpha-1)/p}\right|^p}{(1-r)^{1+\alpha}}\ dr,\ \alpha = \mathrm{const} \in (1,p);$$

$$g_{\alpha,n}(p) = [D_{\alpha,n}(p)]^{-1/p}\,,$$

$$\mu_\alpha(A) := \int_A \frac{dx}{d_\alpha(x)},\ A \subset \Omega;$$

$$\nu_\alpha(B) := \int\int_B \frac{dx\ dy}{|x-y|^{n+\alpha}},\ B \subset \Omega \times \Omega.$$

$$\delta[f](x,y) := f(x) - f(y),\ f : R^n \to R. \tag{4.1}$$

The fractional weight Sobolev's type inequality

$$|f|L_p(R^n, \mu_\alpha) \le g_{\alpha,n}(p)\ |\ \delta[f]\ |L_p(R^n \times R^n, \nu_\alpha) \tag{4.2}$$

was proved by R. L. Frank and R. Seiringer [33] (case $\Omega = R^n_+$) and M.Loss and C.Sloane [39] (general case). See also [32].

**Theorem 4.1.** *Let the function $\delta[f](\cdot,\cdot)$ belongs to some space $G\psi$ on the set $\Omega \times \Omega$ relative the measure $\nu_\alpha$. Put*

$$\theta(p) = g_{\alpha,n}(p) \cdot \psi(p).$$

*Proposition:*

$$||f||G\theta(\Omega, \mu_\alpha) \le 1 \cdot ||\delta[f]||G\psi(\Omega \times \Omega, \nu_\alpha), \tag{4.3}$$

*where the constant "1" in (4.3) is the best possible.*

**Proof** is at the same as in theorem 2.1 and may be omitted.

# 5 Auxiliary results

**Constants** $K(n,s)$.

As long as $\Gamma(\epsilon) \sim 1/\epsilon,\ \epsilon \to 0+$, we deduce at $s \to n-0$

$$K(n,s) \sim \frac{2\pi^{n/2}}{\Gamma(n/2)}\ \frac{1}{n-s}.$$

The point $s = n-0$ is unique point of singularity for the function $s \to K(n,s)$; for instance, $K(n,0) = K(n,0+) = 1$.

**Constants** $D_{\alpha,n}(p)$.

Denote for the values $\alpha = \mathrm{const} > 1$

$$L_\alpha(p) = \int_0^1 \frac{\left|1 - r^{(\alpha-1)/p}\right|^p}{(1-r)^{1+\alpha}}\ dr,\ p \in (\alpha, \infty).$$

Recall that

$$D_{\alpha,n}(p) := 2\pi^{(n-1)/2}\ \frac{\Gamma((1+\alpha)/2)}{\Gamma((n+\alpha)/2)}\ L_\alpha(p).$$

The extreme points $p = \alpha + 0$ and $p \to \infty$ are points of singularity for the function $p \to L_\alpha(p)$.

**A. Case** $p \to \alpha + 0$.

We have:

$$r^{(\alpha-1)/p} = e^{\ln r \cdot (\alpha-1)/p} \sim 1 - \frac{|\ln r|(\alpha-1)}{p},$$

therefore

$$L_\alpha(p) \sim \frac{(\alpha-1)^p}{p^p} \int_0^1 \frac{|\ln r|^p \, dr}{(1-r)^{1+\alpha}} \sim$$

$$\frac{(\alpha-1)^p}{p^p} \int_0^1 (1-r)^{p-1-\alpha} \, dr = \frac{(\alpha-1)^p}{p^p} \cdot (p-\alpha)^{-1}.$$

**B. Case $p \to \infty$.**

We find:

$$L_\alpha(p) \sim \frac{(\alpha-1)^p}{p^p} \int_0^1 \frac{|\ln r|^p \, dr}{(1-r)^{1+\alpha}} \sim$$

$$\frac{(\alpha-1)^p}{p^p} \int_0^1 |\ln r|^p \, dr = \frac{(\alpha-1)^p}{p^p} \cdot \Gamma(p+1).$$

**C. Non-asymptotical approach.**

We will use the following elementary estimate:

$$1 - \sinh(1) \, \epsilon \le e^{-\epsilon} \le 1 - \epsilon, \ \epsilon \in [0,1].$$

Let $\Delta = \mathrm{const} \in (0,1)$; for example, $\Delta = \Delta_0 = 1/2$. We calculate:

$$J := \int_0^1 \frac{|\ln r|^p}{(1-r)^{1+\alpha}} dr = \int_1^\Delta dr + \int_\Delta^1 dr = J_1 + J_2;$$

$$J_1 \le (1-\Delta)^{-1-\alpha} \int_0^\Delta |\ln r|^p \, dr \le$$

$$(1-\Delta)^{-1-\alpha} \int_0^1 |\ln r|^r \, dr = \frac{\Gamma(p+1)}{(1-\Delta)^{1+\alpha})};$$

$$J_2 = \int_\Delta^1 \frac{|\ln r|^p}{(1-r)^{1+\alpha}} dr \le \frac{|\ln \Delta|^p}{((1-\Delta)^p)} \, \frac{1}{p-\alpha},$$

so

$$J \le \frac{\Gamma(p+1)}{(1-\Delta)^{1+\alpha}} + \frac{|\ln \Delta|^p}{((1-\Delta)^p)} \, \frac{1}{p-\alpha},$$

following

$$L_\alpha(p) \le \frac{(\alpha-1)^p}{p^p} \cdot \left[ \frac{\Gamma(p+1)}{(1-\Delta)^{1+\alpha}} + \frac{|\ln \Delta|^p}{((1-\Delta)^p)} \, \frac{1}{p-\alpha} \right].$$

If we choose $\Delta = 1/2$, then

$$L_\alpha(p) \le \frac{(\alpha-1)^p}{p^p} \cdot \left[ 2^{1+\alpha} \Gamma(p+1) + (2 \ln 2)^p \ \frac{1}{p-\alpha} \right],$$

and analogously

$$L_\alpha(p) \ge C^p(\alpha) \cdot \frac{(\alpha-1)^p}{p^p} \cdot \left[ \Gamma(p+1) + \frac{1}{p-\alpha} \right], \ C(\alpha) \in (0,1).$$

**D. Constant of L.Cafarelli, E. Valdinoci, O. Savin.**

There exists a "constant" $Z = Z(n,s,p), \ s \in (0,1), \ p \ge 1$ such that for all measurable set $E \subset R^n$ with positive finite measure $|E|$

$$\int_{R^n \setminus E} \frac{dy}{|x-y|^{n+sp}} \ge Z(n,s,p) \ |E|^{-sp/n},$$

see, e.g. [30], [46]. This constant play very important role in the theory of imbedding of fractional Sobolev's spaces [43].

We will understand as a capacity of the value $Z(n,s,p)$ its maximal value, i.e.

$$Z(n,s,p) \stackrel{def}{=} \inf_{x \in E} \ \inf_{E: |E| \in (0,\infty)} \left\{ \int_{R^n \setminus E} \frac{dy}{|x-y|^{n+sp}} : |E|^{-sp/n} \right\}.$$

Denote also as usually

$$\omega_n = \frac{2 \pi^{n/2}}{\Gamma(n/2)} -$$

the area of surface of unit sphere $R^n$; recall that the volume of unit ball in this space is equal to $\omega_n / n$.

**Proposition 5.1.**

$$(sp)^{-1} \ \omega_n^{1+sp/n} \ n^{-1-sp/n} \le Z(n,s,p) \le (sp)^{-1} \ \omega_n^{1+sp/n} \ n^{-sp/n}.$$

The left - hand side follows immediately from lemma 6.1 in the article [43] after simple calculations; the right - hand side may be obtained by choosing $x = 0$ and $E = \{ y : |y| \le 1. \}$

Obviously, the upper bound in the last inequality is attainable.

**Acknowledgements.**

The authors are grateful to prof. S.V.Astashkin for useful conversations and to Eleonora Di Nezza, Giampiero Palatucci, Enrico Valdinoci for sending their remarkable articles.

# References


[1] Bennet G, Sharpley R. Interpolation of operators. Orlando, Academic Press Inc., (1988).

[2] Capone C., Fiorenza A., Krbec M. On the Extrapolation Blowups in the $L_p$ Scale. Collectanea Mathematica, **48**, 2, (1998), 71-88.

[3] Fiorenza A. Duality and reflexivity in grand Lebesgue spaces. Collectanea Mathematica (electronic version), **51**, 2, (2000), 131-148.

[4] Fiorenza A., and Karadzhov G.E. Grand and small Lebesgue spaces and their analogs. Consiglio Nationale Delle Ricerche, Instituto per le Applicazioni del Calcoto Mauro Picine, Sezione di Napoli, Rapporto tecnico n. 272/03, (2005).

[5] Iwaniec T., and Sbordone C. On the integrability of the Jacobian under minimal hypotheses. Arch. Rat.Mech. Anal., 119, (1992), 129143.

[6] Iwaniec T., P. Koskela P., and Onninen J. Mapping of finite distortion: Monotonicity and Continuity. Invent. Math. 144 (2001), 507-531.

[7] Kozachenko Yu. V., Ostrovsky E.I. (1985). The Banach Spaces of random Variables of subgaussian type. *Theory of Probab. and Math. Stat.* (in Russian). Kiev, KSU, **32**, 43-57.

[8] Liflyand E., Ostrovsky E., Sirota L. Structural Properties of Bilateral Grand Lebesgue Spaces. Turk. J. Math.; **34** (2010), 207-219.

[9] Luigi DOnofrio, Carlo Sbordone, Roberta Schiattarella. Grand Sobolev spaces and their applications in geometric function theory and PDE. Journal of Fixed Point Theory and Applications, June 2013, Volume 13, Issue 2, pp. 309 − 340.

[10] Ostrovsky E.I. Exponential Estimations for Random Fields. Moscow-Obninsk, OINPE, 1999 (Russian).

[11] Ostrovsky E. and Sirota L. Moment Banach spaces: theory and applications. HIAT Journal of Science and Engineering, **C**, Volume 4, Issues 1-2, pp. 233-262, (2007).

[12] E.Ostrovsky, L.Sirota. Adaptive estimation of multidimensional spectral densities. Proceedings of Institute for Advanced Studies, Arad, Israel, (2005), issue 5, p. 42-48.

[13] E. Ostrovsky and L.Sirota. Adaptive multidimensional-time spectral Measurements in technical diagnosis. Communications in dependability and Managements (CDQM), Vol. 9, No 1, (2006), pp. 45-50.

[14] E.Ostrovsky, L.Sirota. *Some singular operators in the bide-side Grang Lebesgue spaces.* All-Russia School-Conference for Undergraduate and Postgraduate Students and Young Scientists "Fundamental Mathematics and its Applications in Natural Sciences", Articles, Mathematics, vol. 2, Ufa: BashSU, (2008), pp. 241-249.

[15] E.Ostrovsky, E.Rogover and L.Sirota. Riesz's and Bessel's operators in in bilateral Grand Lebesgue Spaces. arXiv:0907.3321 [math.FA] 19 Jul 2009.

[16] E. Ostrovsky and L.Sirota. Weight Hardy-Littlewood inequalities for different powers. arXiv:09010.4609v1[math.FA] 29 Oct 2009.

[17] E. Ostrovsky E. Bide-side exponential and moment inequalities for tail of distribution of Polynomial Martingales. Electronic publication, arXiv: math.PR/0406532 v.1 Jun. 2004.

[18] E.Ostrovsky, E.Rogover and L.Sirota. Integral Operators in Bilateral Grand Lebesgue Spaces. arXiv:09012.7601v1 [math.FA] 16 Dez. 2009.

[19] E.Ostrovsky, L.Sirota. Nikolskii-type inequalities for rearrangement invariant spaces. arXiv:0804.2311v1 [math.FA] 15 Apr 2008.

[20] Ostrovsky E., Sirota L. Universal adaptive estimations and confidence intervals in the non-parametrical statistics. arXiv.mathPR/0406535 v1 25 Jun 2004.

[21] Ostrovsky E., Sirota L. Optimal adaptive nonparametric denoising of multidimensional-time signal. arXiv:0809.30211v1 [physics.data-an] 17 Sep 2008.

[22] Ostrovsky E. *Exponential Orlicz's spaces: new norms and applications.*Electronic Publications, arXiv/FA/0406534, v.1, (25.06.2004.)

[23] Ostrovsky E., Sirota L. *Nikolskii-type inequalities for rearrangement Invariant spaces.* Electronic Publications, arXiv:0804.2311 v1 [math.FA] 15 Apr 2008.

[24] Ostrovsky E. Bide-side exponential and moment inequalities for tail of distribution of Polynomial Martingales. Electronic publication, arXiv: math.PR/0406532 v.1 Jun. 2004.

[25] E. Ostrovsky and L. Sirota. *Necessary conditions for fractional Hardy - Sobolew's inequalities.* arXiv:1108.1387v1 [math.FA] 5 Aug 2011

[26] E. Ostrovsky and L. Sirota. *Module of continuity for the functions belonging to the Sobolev-Grand Lebesgue Spaces.* arXiv:1006.4177v1 [math.FA] 21 Jun 2010

[27] E. Ostrovsky and L. Sirota. *Boundedness of operators in bilateral Grand Lebesgue spaces, with exact and weakly exact constant calculation.* arXiv:1104.2963v1 [math.FA] 15 Apr 2011

[28] Talenti G. Inequalities in Rearrangement Invariant Function Spaces. Nonlinear Analysis, Function Spaces and Applications. Prometheus, Prague, **5**, (1995), 177-230.

[29] K. Bogdan and B. Dyda. *The best constant in a fractional Hardy inequality.* Math. Nachr., 284(5-6): 629 − 638, 2011.

[30] L. Caffarelli, E. Valdinoci. *Uniform estimates and limiting arguments for nonlocal minimal surfaces.* Calc. Var. Partial Differential Equations 41 (12), (2011), 203 − 240.

[31] A. Cotsiolis, N.K. Tavoularis. *Best constants for Sobolev inequalities for higher order fractional derivatives.* J. Math. Anal. Appl. 295 (2004), 225 − 236.

[32] Dyda B., Frank R.L. *Fractional Hardy-Sobolev-Maz'ya inequality for domains.* arXiv:1109.6570v1 [math.FA] 29 Sep 2011

[33] R. L. Frank and R. Seiringer. *Sharp fractional Hardy inequalities in half-spaces.* In Around the research of Vladimir Mazya. I, volume 11 of Int. Math. Ser. (N. Y.), pages 161 − 167. Springer, New York, 2010.

[34] G.H.Hardy, J.E. Litlewood and G.Polya. *Inequalities.* Cambridge, (1952).

[35] R. Hurri-Syrjänen and A. V. Vähäkangas. *Characterizations to the fractional Sobolev inequality.* arXiv:1312.3135v1[math.CA] 11Dec 2013.

[36] G. Leoni. *A First Course in Sobolev Spaces.* Grad. Stud. Math., vol. 105, Amer. Math. Soc., Providence, RI, 2009.

[37] E.H.Lieb. *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities.* Ann. of Math., (2), 118 (1983), no 2, 349-374.

[38] E.H. Lieb and M. Loss. *Analysis.* Volume 14 of Graduate Studies in Mathematics, AMS, 1997.

[39] M. Loss and C. Sloane. *Hardy inequalities for fractional integrals on general domains.* J. Funct. Anal., 259(6):13691379, 2010.

[40] M. Ludwig. *Anisotropic fractional Sobolev norms.* Adv. Math., **252**, (2014), 150 − 157.

[41] V. G.Maz'ja. *Sobolev spaces.* Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1985. Translated from the Russian by T. O. Shaposhnikova.

[42] V. Maz'ya, T. Shaposhnikova. *On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces.* J. Funct. Anal. 195 (2002), 230 − 238.

[43] Eleonora Di Nezza, Giampiero Palatucci, Enrico Valdinoci. *Hitchhiker's guide to the fractional Sobolev spaces.* Bull. Sci. math. 136 (2012), 521 − 573.

[44] G. Palatucci, A. Pisante. *Sobolev embeddings and concentration-compactness alternative for fractional Sobolev spaces.* Submitted for publication, available online at http://mipa.unimes.fr/preprints.html.

[45] T. Runst, W. Sickel. *Sobolev Spaces of Fractional Order, Nemytskij Operators, and Nonlinear Partial Differential Equations.* De Gruyter Ser. Nonlinear Anal. Appl., vol. 3, Walter de Gruyter aaand Co., Berlin, 1996.

[46] O. Savin, E. Valdinoci. *Density estimates for a variational model driven by the Gagliardo norm.* arXiv.org/abs/1007.2114v3.

[47] C. A. Sloane. *A fractional Hardy-Sobolev-Mazya inequality on the upper halfspace.* Proc. Amer. Math. Soc., 139:40034016, 2011.

[48] L.N. Slobodetskij. *Generalized Sobolev spaces and their applications to boundary value problems of partial differential equations.* Leningrad. Gos. Ped. Inst. Uch. Zap. 197 (1958), 54 − 112.

[49] Uspenskii S.V. *An imbedding theorem for S.L. Sobolevs classes $W_p^r$ of fractional order.* Dokl. Akad. Nauk SSSR, 130 (1960) 992 993; English translation in: Soviet Math. Dokl. 1 (1960), 132 − 133.

[50] Uspenskii S.V. *Properties of the classes $W_p^r$ with fractional derivatives on differentiable manifold.* Dokl. Akad. Nauk SSSR 132 (1960) 60 62; English translation in: Soviet Math. Dokl. 1 (1960) 495 − 497.